

Erratum

Order and strain in main-chain smectic liquid-crystalline polymers and elastomers

W.H. de Jeu^{1,a}, E.P. Obraztsov¹, B.I. Ostrovskii^{1,2}, W. Ren³, P.J. McMullan³, A.C. Griffin³, A. Sánchez-Ferrer⁴, and H. Finkelmann⁴

¹ FOM Institute for Atomic and Molecular Physics, Kruislaan 407, 1098 SJ Amsterdam, The Netherlands

² Institute of Crystallography, Academy of Sciences of Russia, Leninsky prospect 59, 117333 Moscow, Russia

³ School of Polymer, Textile and Fiber Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA


⁴ Institut für Makromolekulare Chemie, Universität Freiburg, D-79104 Freiburg, Germany

Original article: Eur. Phys. J. E **24**, 399–409 (2007) DOI: 10.1140/epje/i2007-10254-8


Received 28 January 2008

Published online: 1 February 2008 – © EDP Sciences / Società Italiana di Fisica / Springer-Verlag 2008


After the publication of the paper, the authors have remarked a mistake in fig. 2, *i.e.*, in panel (c) one should have Sm-A and not Sm-C. The correct figure is published below.



MC11-pol: K 50°C Sm-C 95°C I
MC11-el: K 41°C Sm-C 110°C I



MeHQ-pol: g -33°C Sm-C 98°C I
MeHQ-el: g -25°C Sm-C 104°C I



TR5-el: Sm-A 52°C I
No glass transition was detected

^a Present address: Polymer Science and Engineering, University of Massachusetts, 120 Governors Drive, Amherst, MA 01003, USA; e-mail: dejeu@mail.pse.umass.edu